

Overview: Cellular Messaging

- Cell-to-cell communication is essential for both multicellular and unicellular organisms
- Biologists have discovered some universal mechanisms of cellular regulation
- Cells most often communicate with each other via chemical signals
- For example, the fight-or-flight response is triggered by a signaling molecule called epinephrine

Figure 11.1




Concept 11.1: External signals are converted to responses within the cell

- Microbes provide a glimpse of the role of cell signaling in the evolution of life

Evolution of Cell Signaling

- The yeast, *Saccharomyces cerevisiae*, have two mating types, **a** and α
- Cells of different mating types locate each other via secreted factors specific to each type
- A **signal transduction pathway** is a series of steps by which a signal on a cell's surface is converted into a specific cellular response
- Signal transduction pathways convert signals on a cell's surface into cellular responses


Figure 11.2



- Pathway similarities suggest that ancestral signaling molecules evolved in prokaryotes and were modified later in eukaryotes
- The concentration of signaling molecules allows bacteria to sense local population density


Figure 11.3

1 Individual rod-shaped cells




0.5 mm

2 Aggregation in progress



3 Spore-forming structure (fruiting body)



2.5 mm

Fruiting bodies






Figure 11.3a



1 Individual rod-shaped cells

© 2011 Pearson Education, Inc.


Figure 11.3b



2 Aggregation in progress

© 2011 Pearson Education, Inc.


0.5 mm



**3 Spore-forming structure
(fruiting body)**


© 2011 Pearson Education, Inc.

Figure 11.3d




Local and Long-Distance Signaling

- Cells in a multicellular organism communicate by chemical messengers
- Animal and plant cells have cell junctions that directly connect the cytoplasm of adjacent cells
- In local signaling, animal cells may communicate by direct contact, or cell-cell recognition




(a) Cell junctions




(b) Cell-cell recognition

- In many other cases, animal cells communicate using **local regulators**, messenger molecules that travel only short distances
- In long-distance signaling, plants and animals use chemicals called **hormones**
- The ability of a cell to respond to a signal depends on whether or not it has a receptor specific to that signal


Figure 11.5



Local signaling




(a) Paracrine signaling



(b) Synaptic signaling

Figure 11.5b



The Three Stages of Cell Signaling:

A Preview

- Earl W. Sutherland discovered how the hormone epinephrine acts on cells
- Sutherland suggested that cells receiving signals went through three processes
 - **Reception**
 - **Transduction**
 - **Response**



Animation: Overview of Cell Signaling

Figure 11.6-1




Figure 11.6-2





Figure 11.6-3



Concept 11.2: Reception: A signaling molecule binds to a receptor protein, causing it to change shape


- The binding between a signal molecule (**ligand**) and receptor is highly specific
- A shape change in a receptor is often the initial transduction of the signal
- Most signal receptors are plasma membrane proteins

Receptors in the Plasma Membrane

- Most water-soluble signal molecules bind to specific sites on receptor proteins that span the plasma membrane
- There are three main types of membrane receptors
 - G protein-coupled receptors
 - Receptor tyrosine kinases
 - Ion channel receptors

- **G-protein-coupled receptor (GPCRs)** are the largest family of cell-surface receptors
- A GPCR is a plasma membrane receptor that works with the help of a **G protein**
- The G protein acts as an on/off switch: If GDP is bound to the G protein, the G protein is inactive

Figure 11.7a



G protein-coupled receptor

© 2011 Pearson Education, Inc.

Figure 11.7b












Figure 11.8




- **Receptor tyrosine kinases (RTKs)** are membrane receptors that attach phosphates to tyrosines
- A receptor tyrosine kinase can trigger multiple signal transduction pathways at once
- Abnormal functioning of RTKs is associated with many types of cancers

Figure 11.7c



- A **ligand-gated ion channel** receptor acts as a gate when the receptor changes shape
- When a signal molecule binds as a ligand to the receptor, the gate allows specific ions, such as Na^+ or Ca^{2+} , through a channel in the receptor

Figure 11.7d



Intracellular Receptors

- Intracellular receptor proteins are found in the cytosol or nucleus of target cells
- Small or hydrophobic chemical messengers can readily cross the membrane and activate receptors
- Examples of hydrophobic messengers are the steroid and thyroid hormones of animals
- An activated hormone-receptor complex can act as a transcription factor, turning on specific genes

Figure 11.9-1




Figure 11.9-2




Figure 11.9-3




Figure 11.9-4





Figure 11.9-5



Concept 11.3: Transduction: Cascades of molecular interactions relay signals from receptors to target molecules in the cell

- Signal transduction usually involves multiple steps
- Multistep pathways can amplify a signal: A few molecules can produce a large cellular response
- Multistep pathways provide more opportunities for coordination and regulation of the cellular response

Signal Transduction Pathways

- The molecules that relay a signal from receptor to response are mostly proteins
- Like falling dominoes, the receptor activates another protein, which activates another, and so on, until the protein producing the response is activated
- At each step, the signal is transduced into a different form, usually a shape change in a protein

Protein Phosphorylation and Dephosphorylation

- In many pathways, the signal is transmitted by a cascade of protein phosphorylations
- **Protein kinases** transfer phosphates from ATP to protein, a process called phosphorylation

- **Protein phosphatases** remove the phosphates from proteins, a process called dephosphorylation
- This phosphorylation and dephosphorylation system acts as a molecular switch, turning activities on and off or up or down, as required

Figure 11.10





Figure 11.10a



Small Molecules and Ions as Second Messengers

- The extracellular signal molecule (ligand) that binds to the receptor is a pathway's “first messenger”
- **Second messengers** are small, nonprotein, water-soluble molecules or ions that spread throughout a cell by diffusion
- Second messengers participate in pathways initiated by GPCRs and RTKs
- Cyclic AMP and calcium ions are common second messengers

Cyclic AMP

- **Cyclic AMP (cAMP)** is one of the most widely used second messengers
- **Adenylyl cyclase**, an enzyme in the plasma membrane, converts ATP to cAMP in response to an extracellular signal

Figure 11.11




Figure 11.11a






Figure 11.11b



- Many signal molecules trigger formation of cAMP
- Other components of cAMP pathways are G proteins, G protein-coupled receptors, and protein kinases
- cAMP usually activates protein kinase A, which phosphorylates various other proteins
- Further regulation of cell metabolism is provided by G-protein systems that inhibit adenylyl cyclase


Figure 11.12



Calcium Ions and Inositol Triphosphate (IP₃)

- Calcium ions (Ca²⁺) act as a second messenger in many pathways
- Calcium is an important second messenger because cells can regulate its concentration

Figure 11.13



Key **High [Ca²⁺]** **Low [Ca²⁺]**

- A signal relayed by a signal transduction pathway may trigger an increase in calcium in the cytosol
- Pathways leading to the release of calcium involve **inositol triphosphate (IP₃)** and **diacylglycerol (DAG)** as additional second messengers



Animation: Signal Transduction Pathways

Figure 11.14-1




Figure 11.14-2





Figure 11.14-3




Concept 11.4: Response: Cell signaling leads to regulation of transcription or cytoplasmic activities

- The cell's response to an extracellular signal is sometimes called the “output response”

Nuclear and Cytoplasmic Responses

- Ultimately, a signal transduction pathway leads to regulation of one or more cellular activities
- The response may occur in the cytoplasm or in the nucleus
- Many signaling pathways regulate the synthesis of enzymes or other proteins, usually by turning genes on or off in the nucleus
- The final activated molecule in the signaling pathway may function as a transcription factor

Figure 11.15



- Other pathways regulate the activity of enzymes rather than their synthesis

Figure 11.16

Reception

Binding of epinephrine to G protein-coupled receptor (1 molecule)



Transduction

Inactive G protein

Active G protein (10^2 molecules)

Inactive adenylyl cyclase

Active adenylyl cyclase (10^2)

ATP

Cyclic AMP (10^4)

Inactive protein kinase A

Active protein kinase A (10^4)

Inactive phosphorylase kinase

Active phosphorylase kinase (10^5)

Inactive glycogen phosphorylase


Active glycogen phosphorylase (10^6)

Response



Glycogen
Glucose 1-phosphate
(10^8 molecules)

- Signaling pathways can also affect the overall behavior of a cell, for example, changes in cell shape

RESULTS



Wild type (with shmoos)

 Δ Fus3 Δ formin

CONCLUSION






Figure 11.17a



Wild type (with shmoos)

© 2011 Pearson Education, Inc.


Figure 11.17b



Δ Fus3

© 2011 Pearson Education, Inc.

Figure 11.17c



Δ formin

© 2011 Pearson Education, Inc.

Fine-Tuning of the Response

- There are four aspects of fine-tuning to consider
 - Amplifying the signal (and thus the response)
 - Specificity of the response
 - Overall efficiency of response, enhanced by scaffolding proteins
 - Termination of the signal

Signal Amplification

- Enzyme cascades amplify the cell's response
- At each step, the number of activated products is much greater than in the preceding step

The Specificity of Cell Signaling and Coordination of the Response

- Different kinds of cells have different collections of proteins
- These different proteins allow cells to detect and respond to different signals
- Even the same signal can have different effects in cells with different proteins and pathways
- Pathway branching and “cross-talk” further help the cell coordinate incoming signals

Figure 11.18




Figure 11.18a







Figure 11.18b



Cell C. Cross-talk occurs between two pathways.




Cell D. Different receptor leads to a different response.

Signaling Efficiency: Scaffolding Proteins and Signaling Complexes

- **Scaffolding proteins** are large relay proteins to which other relay proteins are attached
- Scaffolding proteins can increase the signal transduction efficiency by grouping together different proteins involved in the same pathway
- In some cases, scaffolding proteins may also help activate some of the relay proteins

Figure 11.19




Termination of the Signal

- Inactivation mechanisms are an essential aspect of cell signaling
- If ligand concentration falls, fewer receptors will be bound
- Unbound receptors revert to an inactive state

Concept 11.5: Apoptosis integrates multiple cell-signaling pathways

- **Apoptosis** is programmed or controlled cell suicide
- Components of the cell are chopped up and packaged into vesicles that are digested by scavenger cells
- Apoptosis prevents enzymes from leaking out of a dying cell and damaging neighboring cells

Figure 11.20



© 2011 Pearson Education, Inc.

Apoptosis in the Soil Worm *Caenorhabditis elegans*

- Apoptosis is important in shaping an organism during embryonic development
- The role of apoptosis in embryonic development was studied in *Caenorhabditis elegans*
- In *C. elegans*, apoptosis results when proteins that “accelerate” apoptosis override those that “put the brakes” on apoptosis

Figure 11.21







Figure 11.21a





Apoptotic Pathways and the Signals That Trigger Them

- Caspases are the main proteases (enzymes that cut up proteins) that carry out apoptosis
- Apoptosis can be triggered by
 - An extracellular death-signaling ligand
 - DNA damage in the nucleus
 - Protein misfolding in the endoplasmic reticulum

- Apoptosis evolved early in animal evolution and is essential for the development and maintenance of all animals
- Apoptosis may be involved in some diseases (for example, Parkinson's and Alzheimer's); interference with apoptosis may contribute to some cancers

Figure 11.22





Figure 11.22a

Interdigital tissue



© 2011 Pearson Education, Inc.

Cells undergoing apoptosis





Figure 11.22c



© 2011 Pearson Education, Inc.

Figure 11.UN01




Figure 11.UN02

